

MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS-1963-A

л эі /

. . .

ECULITY CLASSIFICATION OF THIS PAGE (Miten Data F	ritered)		
REPORT DOCUMENTATION PAGE			READ INSTRUCTIONS ORE COMPLETING FORM
None	2 GOVT ACCESSION NO.	3 RECIPIE	NT'S CATALOG NUMBER
TITLE (and Subtitle)		S. TYPE OF	REPORT & PERIOD COVERED
A NOTE ON THE EXTENDED KALMAN FILTER		Final,	11/29/79-11/28/81
		6. PERFOR	MING ORG. REPORT NUMBER
AUTHOR(e)		-	CT OR GRANT NUMBER(4)
Hosam E. Emara-Shabaik		DASG-60-	-80-C-0007
UCLA, School of Engineering & Appli	led Science		
PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRA	M ELEMENT. PROJECT, TASK WORK UNIT NUMBERS
UCLA, School of Engineering and Applied Science Los Angeles, California 90024			None
CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT	DATE
Advanced Technology Center		Marc	h. 1982
U.S. Army Ballistic Missile Defense Command Huntsville, Alabama		13. NUMBER	
MONITORING AGENCY NAME & ADDRESS(II ditterent i	from Controlling Office)		ry CLASS. (of this report) assified
Same as No. 11		15e. DECLA: SCHEDI	SSIFICATION, DOWNGRADING

DISTRIBUTION STATEMENT (of this Report,

Distribution of this document is not limited.

DISTRIBUTION STATEMENT A

Approved for public releases Distribution Unlimited

17 DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)

None

18 SUPPLEMENTARY NOTES

None

19 KEY WORDS (Continue on reverse elde il necessary and identify by block number)

B

20 ABSTRACT (Continue on reverse side if necessary and identify by block number)


See attached

DD , 5000 1473

EDITION OF 1 NOV 65 IS OBSOLETE S/N 0102 LF 014 6601

#### **ABSTRACT**

The extended Kalman Filter (EKF), in the continuous time version, has to be interpreted in the Ito sense. Therefore, processing of the (EKF) algorithm is much more involved than has appeared to be the case. An additional  $\frac{n^2 (n+1)}{2}$  differential equations have to be processed when the EKF is properly interpreted.





Distr	ilution/	
Avai	lability	Codos
Dist	Avail az Specia	od/or
A		

#### A NOTE ON THE EXTENDED KALMAN FILTER

Вy

Hosam E. Emara-Shabaik

March, 1982

Submitted under contract
DASG-60-80-C-0007
Advances In Technology Development
for Exoatmospheric Intercept Systems

U.S. Army Ballistic Missile
Defense Command
Advanced Technology Center
Huntsville, Alabama

School of Engineering and Applied Science University of California Los Angeles, California

## **ABSTRACT**

The extended Kalman Filter (EKF), in the continuous time version, has to be interpreted in the Itô sense. Therefore, processing of the (EKF) algorithm is much more involved than has appeared to be the case. An additional  $\frac{n^2 (n+1)}{2}$  differential equations have to be processed when the EKF is properly interpreted.

#### **PRESENTATION**

Consider the continuous nonlinear stochastic dynamic system and observations given by

$$dx(t) = f(x(t),t) dt + G(t) dw(t), x(t_0) = X_0$$
 (1)

And

$$dz(t) = h(x(t), t) dt + dv(t)$$
 (2)

Where

x(t) & R<sup>n</sup> is an 'n' dimensional state vector.

f(x(t),t) is an 'n' dimensional vector valued function.

 $x_0 \in \mathbb{R}^n$  is an 'n' dimensional Gaussian random vector with mean  $\overline{x}_0$ , and covariance  $P_0$ .

¿(t) & R<sup>m</sup> is an 'm' dimensional observations vector.

h(x(t),t) is an 'm' dimensional vector valued function.

w (t)  $eR^n$  is an 'n' dimensional Wiener process with zero mean and covariance Q(t).

v (t)  $ex^n$  is an 'n' dimensional Wiener process with zero mean and covariance R(t).

Under the assumption that  $x_0$ , w(t), and v(t) are mutually independent of each other, and R(t) considered to be positive definite matrix, the widely used Extended Kalman Filter (EKF) is given by the following set of equations [1]

$$d\hat{x}(t) = f(\hat{x}(t), t) dt + P(t) H^{T}(\hat{x}(t), t) R^{-1}(t) \left[d\hat{x}(t) - h(\hat{x}(t), t) dt\right], \quad \hat{x}(t_{o}) = X_{o}$$
(3)

and

$$\frac{d}{dt} P(t) = F(\hat{x}(t), t) P(t) + P(t)F^{T}(\hat{x}(t), t) + G(t)Q(t)G^{T}(t) - P(t) H^{T}(\hat{x}(t), t) R^{-1}(t) H(\hat{x}(t), t)P(t),$$

$$P(t_{O}) = P_{O} \tag{4}$$

Where

$$H(\hat{x}(t),t) \triangleq \left[ \frac{\partial h_{\hat{x}}(x(t),t)}{\partial x_{\hat{j}}(t)} \Big|_{\hat{x}(t) = \hat{x}(t)} \right]$$
 (5)

and

$$F(\hat{x}(t),t) \triangleq \left[ \frac{\partial f_{L}(x(t),t)}{\partial x_{j}(t)} \Big|_{x(t)=\hat{x}(t)} \right]$$
(6)

It is clear that the matrices  $H(\hat{x}(t),t)$  and  $F(\hat{x}(t),t)$  as given by equations (5) and (6) are state estimate dependent. Hence, also the covariance matrix is state estimate dependent, and a more accurate description of the (EKF) is as follows:

$$d_{\hat{x}}(t) = f(\hat{x}(t), t) dt + K(\hat{x}(t), t) \left[ d_{\hat{z}}(t) - h(\hat{x}(t), t) dt \right],$$

$$\hat{x}(t_0) = \overline{x}_0$$
(7)

 $\frac{d}{dt}P(\hat{x}(t),t) = F(\hat{x}(t),t)P(\hat{x}(t),t) + P(\hat{x}(t),t)F^{T}(\hat{x}(t),t) + G(t)Q(t)G^{T}(t) -$ 

$$P(\hat{x}(t),t)H^{T}(\hat{x}(t),t)R^{-1}(t)H(\hat{x}(t),t)P(\hat{x}(t),t)P(t_{o})=P_{o}$$
 (8)

Where

 $K(\hat{x}(t),t)=P(\hat{x}(t),t)H^{T}(\hat{x}(t),t)R^{-1}(t)$  is the filter's gain matrix.

Now, it is easy to recognize that equation (7) is an Itô stochastic differential equation. Hence, it should be integrated in the Itô sense. Therefore, whenever numerical simulations of the (EKF) are carried out, equation (7) has to be transformed into an equivalent differential equation in an ordinary sense such as the Stratonovich sense. The Stratonovich equivalent of (7) is obtained by modifying the ith component

 $f_i(\hat{x}(t),t)$  by the following quantity

$$\frac{1}{2} \sum_{\ell=1}^{n} \sum_{j=1}^{m} K_{\ell j} (\hat{\hat{x}}(t), t) \frac{\partial K_{ij}(\hat{\hat{x}}(t), t)}{\partial \hat{x}_{\ell}(t)}.$$

Therefore, for the completeness of the EKF, the expression

$$\frac{\partial K_{ij}(\hat{x}(t),t)}{\partial \hat{x}_{i}(t)}$$
 for all i, j, and 2

has to be included. Denote  $\frac{\partial K(\hat{x}(t),t)}{\partial \hat{x}_{i}(t)}$  by

$$\hat{\mathbf{x}}_{i}$$
  $(\hat{\mathbf{x}}(t),t)$ , and  $\frac{\partial P(\hat{\mathbf{x}}(t),t)}{\partial \hat{\mathbf{x}}_{i}}$  by  $\hat{\mathbf{x}}_{i}$   $(\hat{\mathbf{x}}(t),t)$ , then

$$K_{\hat{\mathbf{x}}_{\hat{\mathbf{i}}}}^{\hat{\mathbf{x}}}(\hat{\mathbf{x}}(t),t) = \left[P(\hat{\mathbf{x}}(t),t)\frac{\partial \mathbf{H}^{T}(\hat{\mathbf{x}}(t),t)}{\partial \hat{\mathbf{x}}_{\hat{\mathbf{i}}}} + \right]$$

$$\frac{\partial P(\hat{x}(t),t)}{\partial \hat{x}_{i}} H^{T}(\hat{x}(t),t) \right] R^{-1}(t)$$
(9)

And for  $P_{\hat{x}_i}$   $(\hat{x}(t),t)$ , assuming that  $P(\hat{x}(t),t)$ 

is continuous in both arguments, we have

$$\frac{d}{dt} \hat{P}_{\hat{x}_{i}} (\hat{\hat{x}}(t),t) = F(\hat{\hat{x}}(t),t) P_{\hat{x}_{i}} (\hat{\hat{x}}(t),t) + F_{\hat{x}_{i}} (\hat{\hat{x}}(t),t) P(\hat{\hat{x}}(t),t) + P(\hat{\hat{x}}(t),t) P_{\hat{x}_{i}} (\hat{\hat{x}}(t),t) P(\hat{\hat{x}}(t),t) + P(\hat{\hat{x}}(t),t) P_{\hat{x}_{i}} (\hat{\hat{x}}(t),t) P(\hat{\hat{x}}(t),t) P(\hat{x}(t),t) P(\hat{x}(t),t) P(\hat{x}(t),t) P(\hat{x}(t),t) P(\hat{x}(t),t) P(\hat{x}(t),t$$

$$- P_{\hat{x}_{i}} (\hat{x}(t), t) H^{T} (\hat{x}(t), t) R^{-1}(t) H(\hat{x}(t), t) P(\hat{x}(t), t)$$

$$-P(\hat{x}(t),t)H_{\hat{x}_{i}}^{T}(\hat{x}(t),t)R^{-1}(t)H(\hat{x}(t),t)P(\hat{x}(t),t)$$

- 
$$P(\hat{x}(t),t)H^{T}(\hat{x}(t),t) R^{-1}(t) H_{\hat{x}_{i}}(\hat{x}(t),t) P(\hat{x}(t),t)$$

$$- P(\hat{\hat{x}}(t),t) H^{T}(\hat{\hat{x}}(t),t) R^{-1}(t) H(\hat{\hat{x}}(t),t) P_{\hat{x}_{\hat{1}}} (\hat{\hat{x}}(t),t),$$

$$\hat{\mathbf{x}}_{i} \quad (\mathbf{t}_{o}) = 0 \tag{10}$$

for every i = 1, 2, ....,n

where

$$H_{\hat{\mathbf{X}}_{\underline{i}}}(\hat{\underline{x}}(t),t) = \frac{\partial H(\hat{\underline{x}}(t),t)}{\partial \hat{x}_{\underline{i}}} \triangleq \frac{\partial}{\partial \hat{x}_{\underline{i}}} \left[ \frac{\partial h_{\underline{t}}(\underline{x}(t),t)}{\partial x_{\underline{j}}} \Big|_{\underline{x}(t) = \hat{\underline{x}}(t)} \right]$$

$$= \left[ \frac{\partial^{2} h_{i}(\underline{x}(t), t)}{\partial x_{i} \partial x_{j}} \Big|_{\underline{x}(t) = \hat{\underline{x}}(t)} \right]$$
(11)

and

$$F_{\hat{\mathbf{x}}_{\mathbf{i}}}(\hat{\underline{\mathbf{x}}}(t),t) = \frac{\partial F(\hat{\underline{\mathbf{x}}}(t),t)}{\partial \hat{\mathbf{x}}_{\mathbf{i}}} \stackrel{\underline{\Delta}}{=} \frac{\partial}{\partial \hat{\mathbf{x}}_{\mathbf{i}}} \left[ \frac{\partial f_{\underline{\mathbf{t}}}(\underline{\mathbf{x}}(t),t)}{\partial \mathbf{x}_{\mathbf{j}}} \Big|_{\hat{\underline{\mathbf{x}}}(t) = \underline{\mathbf{x}}(t)} \right]$$

$$= \left[ \frac{\partial^{2} f_{\underline{\mathbf{t}}}(\underline{\mathbf{x}}(t),t)}{\partial \mathbf{x}_{\mathbf{i}} \partial \mathbf{x}_{\mathbf{j}}} \Big|_{\underline{\mathbf{x}}(t) = \hat{\underline{\mathbf{x}}}(t)} \right]. \tag{12}$$

### Conclusion:

The continuous time EKF has to be interpreted in the Ito Therefore, the actual computational requirement for implementing the EKF is increased. An extra n matrix differential equations for  $P_{\hat{x}_i}$  ( $\hat{x}(t)$ ,t) , i=1,2,...n are coupled to the differential matrix Ricatti equation for  $P(\hat{x}(t),t)$ . Also, the nonlinearities f(x(t),t) and h(x(t),t)are assumed to have mixed second order partial derivatives with respect to the components of x(t). Of course, if an integration routine that conforms with the Ito definition of stochastic integral is used in processing the (EKF) algorithm, then it is sufficient to process equations (7) and (8). Among the widely used integration routines, the rectangular routine is the only one that conforms with the Ito definition of the stochastic integral. The discussion in this note applies in a similar way to any of the many suboptimal filters whose gain and, or covariance is estimate dependent.

In practical implementation, the filter equations (3) and (4) are discretized. Usually the time increments are taken to be small enough to assure acceptable accuracy. In order to conform with itô's definition of stochastic integral, the system matrices  $F(\hat{x}(t),t)$  and  $H(\hat{x}(t),t)$  should be held at their values corresponding to the beginning of each particular increment.

# References

200

A. H. Jazwinski, Stochastic Processes And Filtering
Theory, Academic Press, New York, 1970.

DATE FILMED

DTIC